

AWARD NUMBER: W81XWH-15-1-0345

TITLE: A Novel Animal Model for Investigating the Neural Basis of Focal Dystonia

PRINCIPAL INVESTIGATOR: Evinger, Leslie

CONTRACTING ORGANIZATION: The Research Foundation of State University
Stony Brook, NY 11794

REPORT DATE: September 2016

TYPE OF REPORT: Annual Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

*Form Approved
OMB No. 0704-0188*

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE September 2016			2. REPORT TYPE Annual		3. DATES COVERED 1 Sep 2015 – 31 Aug 2016
A Novel Animal Model for Investigating the Neural Basis of Focal Dystonia			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER W81XWH-15-1-0345		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Leslie Evinger			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Research Foundation State University of New York Office of Sponsored Programs Stony Brook University Stony Brook, NY 11794-3362			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The overall goal of the project was to develop an animal model of the focal dystonia benign essential blepharospasm. Consistent with the widely held view that dystonia results from an interaction between a predisposing condition and an environmental trigger, we proposed to use 7 Hz deep brain stimulation of the basal ganglia as the predisposing condition and dry eye as an environmental trigger. Based on experiments during the 1 st year of the grant, our publication, Evinger, J. Neuro-Ophthalmol. 35:374-379, 2015 demonstrated the feasibility of this approach. In addition, we began important control experiments and preliminary recordings from the superior colliculus.					
15. SUBJECT TERMS Dystonia, benign essential blepharospasm, dry eye, motor plasticity, superior colliculus					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Unclassified	18. NUMBER OF PAGES 14	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
1. Introduction.....	4
2. Keywords.....	4
3. Accomplishments.....	4
4. Impact.....	7
5. Changes/Problems.....	7
6. Products.....	7
7. Participants & Other Collaborating Organizations.....	9
8. Special Reporting Requirements.....	
9. Appendices.....	10

1. Introduction

The focal dystonia benign essential blepharospasm (BEB), arises from the convergence of a predisposing condition and an environmental trigger (Hallett et al., 2008). The overall goal of our project was to develop an animal model of BEB based on the hypothesis that hypersynchronized, 7 Hz neuronal oscillations of the basal ganglia created the predisposing condition and that eye irritation from dry eye was the environmental trigger. The basis for our hypothesis that hypersynchronized basal ganglia activity creates a predisposing condition is our demonstration that hypersynchronized oscillations in the basal ganglia produced by 7 Hz deep brain stimulation of the subthalamic nucleus (**STN DBS**) exaggerated neural plasticity in normal rats (Kaminer et al., 2014). Knowing that neural plasticity enables the brain to modify blinking to compensate for dry eye (Evinger et al., 2002; Schicatano et al., 2002), we predicted that combining 7 Hz DBS and dry eye would exaggerate neural plasticity and force the normally compensatory adaptive processes in response to dry eye to transform into the characteristics of BEB, spasms of lid closure, excessive blinking, and trigeminal hyperexcitability. We proposed two Specific Aims to test our hypothesis. The first Specific Aim was to demonstrate that synchronized theta oscillations in the basal ganglia exaggerate plasticity in the cerebellum and the excitability of trigeminal blink circuits as occurs in BEB patients. The Major Tasks to accomplish Specific Aim 1 were: 1) to investigate effects of synchronized basal ganglia oscillations on activity of the deep cerebellar nucleus neurons; and 2) to investigate the effects of synchronized basal ganglia oscillations on activity of the superior colliculus neurons. The second Specific Aim was to demonstrate that synchronized theta (7 Hz) oscillations established in the basal ganglia are sufficient to predispose mammals to develop BEB. The Major tasks to accomplish Specific Aim 2 were: 1) to determine whether combining synchronized basal ganglia theta oscillations with corneal irritation is sufficient to develop spasms of lid closure and other characteristics of the focal dystonia BEB; and 2) to perform control experiments to determine that theta frequency is critical in enabling the development of spasms of lid closure. We completed Major Task 1 of Specific Aim 2 and acquired significant preliminary data for Major Task 2 of Specific Aim 2 in the first year of the award.

2. Keywords

Dystonia, benign essential blepharospasm, dry eye, motor plasticity, basal ganglia, deep brain stimulation

3. Accomplishments

Major goals of the project

The overarching goal of the investigations proposed in this grant was to test our hypothesis that in response to dry eye, the hypersynchronized theta oscillations in the basal ganglia exaggerated neural plasticity of predisposed individuals to develop the focal dystonia benign essential blepharospasm (BEB). We proposed that in the presence of the predisposing condition the normally compensatory adaptive processes engendered by dry eye transformed into the characteristic spasms of lid closure, excessive blinking, and trigeminal hyperexcitability characteristic of BEB. To test this hypothesis, the 1st Specific Aim of the project was to demonstrate that synchronized theta oscillations in the basal ganglia exaggerate plasticity in the cerebellum and excitability of trigeminal blink circuits. Major Task 1 of Specific Aim 1 was to investigate effects of synchronized basal ganglia oscillations on activity of the deep cerebellar nucleus neurons. Major Task 2 of Specific Aim 1 was to investigate the effects of synchronized basal ganglia oscillations on the activity of the superior colliculus neurons. The 2nd Specific Aim of the project was to demonstrate that synchronized theta oscillations established in the basal ganglia are sufficient to predispose mammals to develop blepharospasm. Major Task 1 of Specific Aim 2 was to determine whether combining synchronized basal ganglia theta oscillations with corneal irritation is sufficient to

develop spasms of lid closure and other characteristics of the focal dystonia BEB. Major Task 2 of Specific Aim 2 was to perform control experiments to determine that theta frequency is critical in enabling the development of spasms of lid closure.




Fig. 1. An animal model of benign essential blepharospasm using 7 Hz deep brain stimulation (DBS) as the predisposing condition. A. A recording of spasms of lid closure and excessive blinking by a rat with dry eye receiving 7 Hz subthalamic nucleus (STN) DBS. B. Average number of blinks (# Blinks), blink amplitude (Blink amp), blink duration (Blink Dur), number of spasms (# Spasms), amplitude of spasms (Spasm Amp), and duration of spasms (Spasm Dur) relative to 7 Hz STN DBS alone condition. Spasms were lid closures lasting >100 milliseconds. Error bars are SEM. *P < 0.05; ***P < 0.001. from: Evinger J. *Neuro-Ophthalmology*, 2015;35:374–379

ganglia. Rats in this paradigm developed spasms of lid closure and excessive blinking (Fig. 1 taken from Appendix). These data demonstrated that the theta frequency basal ganglia hypersynchronization created by STN DBS interacted with the plastic compensatory processes initiated by dry eye to create spasms of lid closure typical of BEB. This result supported our hypothesis that theta frequency STN DBS exaggerates the plasticity initiated by dry eye to create spasms of lid closure typical of BEB.

Specific Aim 2, Major Task 2: As described in the project narrative of the grant proposal (page 11), an important control experiment was to demonstrate that the 7 Hz STN DBS is the critical frequency to exaggerate blink plasticity in dry eye into spasms of lid closure. 130 Hz STN DBS does not affect blink plasticity in normal rats (Kaminer et al., 2014), but the appropriate comparison is the dry eye combined with 130 Hz STN DBS because dry eye initiates blink plasticity to compensate for the eye irritation of dry eye. With dry eye initiated plasticity, any frequency of STN DBS might affect blink plasticity. Our hypothesis predicted that the combination of dry eye and 130 Hz STN DBS should not exaggerate blink plasticity. As described on page 8 of the grant proposal project narrative (Trigeminal Reflex Blink Gain Paradigm), we employed our previously published paradigm that decreased trigeminal reflex blink gain in rodents as well as humans (Mao and Evinger, 2001; Ryan et al., 2014) to investigate blink plasticity. In this paradigm, high frequency stimulation of the supraorbital branch of the trigeminal nerve reduced the amplitude of subsequent reflex blinks, a decrease in blink gain. Our data demonstrated that 7 Hz STN DBS (Fig. 2, red) significantly exaggerated the reduction in blink gain relative to normal rats with No STN DBS (Fig. 2, blue) and rats with dry eye receiving 130 Hz STN DBS (Fig. 2, purple). Importantly, there was no




Fig 2. Comparison of blink plasticity among normal rats with 7 Hz DBS and no DBS and rats with dry eye receiving 130 Hz DBS. Error bars are SEM.

What was accomplished under these goals?

In the first year, we completed Major Task 1 and collected preliminary data on Major Task 2 of Specific Aim 2.

Specific Aim 2, Major Task 1: We combined dry eye with theta, 7 Hz, deep brain stimulation (STN DBS) of the subthalamic nucleus of the basal

ganglia. Rats in this paradigm developed spasms of lid closure and excessive blinking (Fig. 1 taken from Appendix).

These data demonstrated that the theta frequency basal ganglia hypersynchronization created by STN DBS interacted with the plastic compensatory processes initiated by dry eye to create spasms of lid closure typical of BEB. This result supported our hypothesis that theta frequency STN DBS exaggerates the plasticity initiated by dry eye to create spasms of lid closure typical of BEB.

With dry eye initiated plasticity, any frequency of STN DBS might affect blink plasticity. Our hypothesis predicted that the combination of dry eye and 130 Hz STN DBS should not exaggerate blink plasticity. As described on page 8 of the grant proposal project narrative (Trigeminal Reflex Blink Gain Paradigm), we employed our previously published paradigm that decreased trigeminal reflex blink gain in rodents as well as humans (Mao and Evinger, 2001; Ryan et al., 2014) to investigate blink plasticity. In this paradigm, high frequency stimulation of the supraorbital branch of the trigeminal nerve reduced the amplitude of subsequent reflex blinks, a decrease in blink gain. Our data demonstrated that 7 Hz STN DBS (Fig. 2, red) significantly exaggerated the reduction in blink gain relative to normal rats with No STN DBS (Fig. 2, blue) and rats with dry eye receiving 130 Hz STN DBS (Fig. 2, purple). Importantly, there was no

significant difference in the plasticity exhibited by normal rats without STN DBS and rats with dry eye receiving 130 Hz STN DBS. Moreover, the rats with dry eye and 130 Hz STN DBS did not develop spasms of lid closure as occurs with 7 Hz STN DBS. Thus consistent with our hypothesis, the data demonstrated that combining dry eye with high frequency STN DBS did not exaggerate blink plasticity. The second control experiment of Major Task 2, combining dry eye with 16 Hz STN DBS is ongoing.

Changes to SOW Order

The decision to begin our investigations with Specific Aim 2 instead of Specific Aim 1 resulted from the lack of personnel available to work on the experiments in Specific Aim 1. The microelectrode recording experiments of Specific Aim 1 require assistance from an individual with some experience in single neuron recording. I was unable to hire a postdoctoral fellow in the first year and have been unable to attract a graduate student to replace the one who just graduated. In contrast, the experiments of Specific Aim 2 could be performed by closely supervised undergraduate students currently in my laboratory. To enable me to start the experiments of Specific Aim 1, I hired a technician who can assist me with the microelectrode recording experiments. We have implanted microelectrode arrays and DBS electrodes in rats and will begin the experiments of Specific Aim 1 in January 2017. We expect to complete the experiments in Major Task 2 of Specific Aim 1 and begin the experiments of Major Task 1 of Specific Aim 1 by the time for the 2017 renewal. The choice of beginning with Major Task 2 rather than Major Task 1 of Specific Aim 1 was that the relative ease of recording from superior colliculus compared to the cerebellar interpositus nucleus enables me to bring the research technician up to speed more quickly.

What opportunities for training and professional development has the project provided?

The graduate student had the opportunity to acquire technical skills in animal behavior, surgical procedures, and alert rodent brain recording.

How were the results disseminated to communities of interest?

We published one manuscript.




Fig. 3. Superior colliculus LFP associated with blinks. A) Averaged LFP (blue) and rectified reflex blink EMG (red). B) Averaged LFP and blink EMG synchronized to occurrence of a spontaneous blink.

What are the plans for the next reporting period to accomplish the goals?

The work during the next reporting period will focus on completing the control experiments of Major Task 2 of Specific Aim 2 as well as the experiments of Major Task 2 of Specific Aim 1. We also hope to begin the experiments of Major Task 1 of Specific Aim 1.

Major Task 2 of Specific Aim 1 (Project Narrative page 9): The goal of these experiments is to determine how different frequencies of basal ganglia oscillations modify superior colliculus activity and to correlate these changes with alterations in trigeminal reflex blink excitability, a primary characteristic of BEB. We will record from the intermediate and deep layers of the superior colliculus receiving spinal trigeminal inputs. Our data demonstrate that both 7 and 16 Hz STN DBS increase reflex blink excitability and that 130 Hz STN DBS has no effect (Kaminer et al., 2014). We will compare the activity of superior colliculus neurons and local field potentials (LFP) with all three frequencies to the No DBS condition. Each day, rats will receive 7 Hz STN DBS, 16 Hz STN DBS, 130 Hz STN DBS, and No DBS blocks of thirty trials of pairs of 2T SO stimuli with a 100 ms interstimulus interval

every 20 ± 5 s. These experiments are already in progress (Fig. 3).

Major Task 1 of Specific Aim 1 (Project Narrative page 9):

The goal of these experiments is to determine how different frequencies of basal ganglia oscillations modify cerebellar interpositus (IP) activity and to correlate these changes with shifts in blink plasticity. As our data demonstrate that 7 Hz STN DBS exaggerates blink plasticity, 16 Hz STN DBS impairs blink plasticity, and 130 Hz STN DBS has no effect on the blink plasticity of normal rats (Kaminer et al., 2014), we will compare IP neural activity across all three frequencies. We will simultaneously record unitary activity and local field potentials (LFP) from blink related IP regions during SO stimulation before, during, and after our trigeminal reflex blink gain paradigm (page 8, project narrative). We will compare the activity of individual IP neurons and LFP before the trigeminal reflex blink gain paradigm to their activity after the trigeminal reflex blink gain paradigm.

4. Impact

What was the impact on the development of the principal discipline(s) of the project?

The preliminary data had an impact on the interpretation of the Neuro-Ophthalmology community as can be seen in the editorial on our manuscript in the Appendix (KB Digre, "Benign Essential Blepharospasm—There Is More to It Than Just Blinking, J. Neuro-Ophthalmol. 35(4):379-381, 2015).

What was the impact on other disciplines?

Nothing to Report

What was the impact on technology transfer?

Nothing to Report

What was the impact on society beyond science and technology?

Nothing to Report

5. Changes/Problems

Changes in approach and reasons for change

There were no changes in approach.

Actual or anticipated problems or delays and actions or plans to resolve them

The one delay in the work has been the difficulty of finding a postdoctoral fellow to participate in the project. I made offers to two individuals. For personal reasons, these two decided not to come to Stony Brook University. My search for a postdoctoral fellow continues. I have also been unable to attract a new graduate student to replace the one who graduated. To speed up the research, I have hired a new research technician to assist with the microelectrode studies of Specific Aim 1.

Changes that had a significant impact on expenditures

Nothing to Report

Significant changes in use or care of vertebrate animals

Nothing to Report

6. Products

Journal publications

"Benign Essential Blepharospasm is a Disorder of Neuroplasticity: Lessons from Animal Models." J. Neuro-ophthalmol. 35:374-379, 2015

Books or other non-periodical, one time publications

None to Report

Other publications, conference papers, and presentations

International:

"Trying to Raise the Window Shades: The Functional Blindness of Benign Essential Blepharospasm"

Invited lecture at Cardiff University, October 28, 2015

Website(s) or other internet site(s)

None to Report

Technologies or techniques

None to Report

Inventions, patent applications, and/or licenses

None to Report

Other Products

None to Report

7. Participants & other collaborating organizations

What individuals have worked on the project?

Name:	Leslie Craig Evinger	Patricia Enmore	Ashley Culoso	Donna Schmidt
Project Role:	PI	Graduate Student	Technician	Technician
Research Identifier:	0000-0002-0039-3348			
Nearest Person Month Worked:	12	12	3	6
Contribution:	Experimental design, manuscript preparation, performing experiments	Performing experiments	Performing experiments	Lab manager, histology
Funding Support:				

Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period?

None to Report

What other organizations were involved as partners?

None to Report

8. Special reporting requirements

Not Applicable

9. Appendices

"Benign Essential Blepharospasm is a Disorder of Neuroplasticity: Lessons from Animal Models." J. Neuro-ophthalmol. 35:374-379, 2015

References

- Evinger C, Mao JB, Powers AS, Kassem IS, Schicatano EJ, Henriquez VM, Peshori KR (2002) Dry eye, blinking, and blepharospasm. Mov Disord 17 Suppl 2:S75-78.
- Hallett M, Evinger C, Jankovic J, Stacy M (2008) Update on blepharospasm: Report from the BEBRF International Workshop. Neurology 71:1275-1282.
- Kaminer J, Thakur P, Evinger C (2014) Frequency matters: beta-band subthalamic nucleus deep-brain stimulation induces Parkinsonian-like blink abnormalities in normal rats. Eur J Neurosci 40:3237-3242.
- Mao JB, Evinger C (2001) Long-term potentiation of the human blink reflex. J Neurosci 21:RC151.
- Ryan M, Kaminer J, Enmore P, Evinger C (2014) Trigeminal high-frequency stimulation produces short- and long-term modification of reflex blink gain. J Neurophysiol 111:888-895.
- Schicatano EJ, Mantzouranis J, Peshori KR, Partin J, Evinger C (2002) Lid restraint evokes two types of motor adaptation. J Neurosci 22:569-576.

Benign Essential Blepharospasm is a Disorder of Neuroplasticity: Lessons From Animal Models

Craig Evinger, PhD

Journal of Neuro-Ophthalmology 2015;35:374–379
doi: 10.1097/WNO.0000000000000317
© 2015 by North American Neuro-Ophthalmology Society

Effectively modeling benign essential blepharospasm (BEB) requires mimicking its root causes. Current evidence points to BEB arising from the confluence of a genetic predisposing condition and an environmental trigger (1). In this “2 hit” hypothesis, the appropriate environmental trigger engenders dystonic behavior because the predisposing condition creates inappropriate brain functioning. Epidemiological studies demonstrate that eye irritation from dry eye, blepharitis, or keratoconjunctivitis is the environmental trigger (1–6). The strength of the association between dry eye and BEB increases in the fifth and sixth decades of life (6) when BEB typically arises (7). Available data strongly support that the predisposing condition is genetic (1,8–12). There is evidence for an autosomal-dominant gene with reduced penetrance contributing to BEB (9,13), but current studies fail to identify any specific genes (8,14). Thus, creating a useful animal model of BEB must involve combining an environmental trigger with a predisposing condition.

Another goal of an animal model is to reproduce the typical symptoms of BEB. The hallmark of BEB is excessive involuntary bilateral lid closure primarily involving the orbicularis oculi muscles (1,15–18). In addition to lid spasms, patients with BEB exhibit trigeminal hyperexcitability (1,15,19–22), an elevated spontaneous blink rate (23), and photophobia (1,24–26). These characteristics are consistent with eye irritation serving as the environmental trigger for BEB because they all appear in patients with dry eye (21,27,28). This relationship between eye irritation and BEB

characteristics indicates that eye irritation should be 1 component of an animal model and that the predisposing condition should cause the adaptive changes in eyelid control in response to dry eye to develop into BEB-like characteristics.

Current evidence demonstrates that trigeminal blink circuits undergo plastic, adaptive modifications to compensate for the rapid breakup of the corneal tear film in dry eye (29–32). Dry eye or eye irritation elevates trigeminal blink amplitude and duration to increase meibomian gland secretion and enhance restoration of the tear film (20,32–37). Blink frequency increases to reform the tear film more regularly (20,36–40). The trigeminal reflex blink circuit becomes hyperexcitable to allow tear film breakup to evoke a reflex blink more readily (20,21,32). Finally, the trigeminal reflex blink circuit responds to a single reflex evoking stimulus with multiple blinks to help restore the tear film (20,21,32).

A simple experiment demonstrates that these modifications are part of a compensatory plastic change occurring in the trigeminal complex (32). Within 30 minutes of restraining 1 eyelid to make blinking more difficult, stimulating the supraorbital nerve ipsilateral to the restrained eyelid evokes hyperexcitable reflex blinks and additional blinks in both eyelids. Stimulating the supraorbital nerve contralateral to the restrained eyelid, however, elicits normal blinks in both eyelids. This pattern would occur only if the trigeminal complex receiving signals of corneal irritation from eyelid restraint expressed the plastic changes. Thus, eye irritation initiates plastic compensatory changes in blinking that could be exaggerated in BEB to produce the eyelid abnormalities of this focal dystonia.

We hypothesize that the predisposing condition exaggerates neuroplasticity so that modifications in response to eye irritation become maladaptive and amplify into the characteristics of BEB. There is significant evidence for exaggerated plasticity in dystonia (41,42). With generalized dystonia, homeostatic synaptic plasticity in the striatum is abnormal (43,44). Exaggerated associative plasticity accompanies focal hand dystonia (45–48). Important for our hypothesis, exaggerated plasticity of the trigeminal blink reflex accompanies BEB (49).

Department Neurobiology and Behavior and Ophthalmology, SUNY Stony Brook, Stony Brook, New York.

Supported by grants from NIH (EY07391) and the Thomas Hartman Parkinson Research Center.

The author reports no conflicts of interest.

Address correspondence to Craig Evinger, PhD, Department Neurobiology and Behavior and Ophthalmology, SUNY Stony Brook, Stony Brook, NY 11790-5230; E-mail: leslie.evinger@stonybrook.edu

Our initial rodent model of BEB (50) used a small reduction of substantia nigra dopamine neurons to create the predisposing condition and crushing 1 branch of the facial nerve innervating the orbicularis oculi to generate the environmental trigger. The choice of dopamine depletion as a predisposing condition came from observations showing that baboons undergoing poisoning with the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) of dopamine neurons exhibited dystonia before developing Parkinsonian movement abnormalities (51) and that there was a disruption of D₂ receptors in patients with BEB (52,53). Thus, changes in dopamine levels or the functioning of specific dopamine receptor subtypes could create the “predisposing condition” for BEB. For an environmental trigger, we created a transient eye irritation by crushing a branch of the facial nerve that provides approximately 30% of the orbicularis oculi innervation. This procedure produced a transient dry eye condition because the weakened eyelid became less effective at restoring the tear film with each blink. The condition was only temporary, however, because regeneration of the crushed nerve branch restored complete lid function within 3 weeks.

In the Schicatano model (50), the BEB-like spasms of lid closure only occurred with the combination of the environmental trigger and the dopaminergic predisposing condition. In the absence of the predisposing condition, the environmental trigger of transient eye irritation slightly increased trigeminal reflex blink excitability and resulted in the development of additional blinks similar to those seen in human dry eye (20,21). Without the environmental trigger, the predisposing condition of a small dopamine neuron loss slightly increased trigeminal reflex blink excitability but did not generate spasms of lid closure. Combining the predisposing condition and the environmental trigger, however, caused long-lasting spasms of lid closure, dramatically elevated trigeminal reflex blink excitability, and increased spontaneous blinking similar to the pattern of blink abnormalities of patients with BEB. These BEB-like characteristics continued after the facial nerve regained full function and eliminated the dry eye. Thus, the BEB-like characteristics of this animal model seemed to result from an exaggeration of the normally compensatory process evoked by eye irritation.

The Schicatano BEB model also was consistent with the important interactions between the cerebellum and basal ganglia that underlie dystonia (54–63). Previous studies demonstrated that the cerebellum was essential for adaptive responses to the eye irritation created by eyelid restraint. Lesions of the cerebellum (30,31) blocked the increases in blink amplitude and duration initiated by eye irritation (20,32–37). Recordings from blink-related neurons in the cerebellar interpositus nucleus revealed the changes in cerebellar activity that accounted for the changes in blink amplitude and duration associated with lid restraint (29). Although the Schicatano model supported the 2 hit hypothesis as the basis of BEB and identified the basal ganglia and cerebellum as key players in this focal dystonia, the model did not explain how the predisposing condition created the

exaggerated plasticity that allowed normally adaptive modification to eye irritation to swell into spasms of lid closure.

We hypothesize that the key to the exaggerated plasticity of dystonia is hypersynchronized low-frequency oscillations of basal ganglia activity. Basal ganglia neurons in patients with Parkinson disease and animal models of Parkinson disease exhibit hypersynchronized oscillations in the broad beta band, 10–30 Hz (64–71). In contrast, basal ganglia neurons in dystonic patients exhibit hypersynchronized oscillations in the theta band, 3–10 Hz (71–74). Although the role of these oscillations in modifying voluntary movement is unclear (66,73,75–81), our study in rodents demonstrate that these basal ganglia oscillations modify trigeminal reflex blink plasticity (82).

We directly tested the role of basal ganglia oscillations in blink plasticity by delivering deep brain stimulation to the basal ganglia subthalamic nucleus of normal rats undergoing a blink plasticity paradigm (82). The procedure was a cerebellar-dependent plasticity paradigm that we developed for humans (83) and modified for rodents (84). Other investigators used this paradigm to demonstrate impaired blink plasticity with Parkinson disease (85), but exaggerated blink plasticity with BEB (49). If the frequency of basal ganglia oscillations modulates brainstem plasticity, then beta frequency deep brain stimulation in normal rats should impair trigeminal reflex blink plasticity, whereas theta frequency, 7 Hz, deep brain stimulation should exaggerate blink plasticity. The Kaminer et al study (82) demonstrated the validity of this postulation. Beta frequency, 16 Hz, deep brain stimulation impaired blink plasticity, whereas theta frequency, 7 Hz, deep brain stimulation exaggerated trigeminal reflex blink plasticity in normal rats. Deep brain stimulation at 130 Hz, a therapeutic frequency for deep brain stimulation in humans (86), however, did not affect blink plasticity in normal rats. Thus, hypersynchronized theta frequency basal ganglia oscillations could create a predisposing condition in which adaptive plasticity initiated by eye irritation exaggerated into spasms of lid closure typical of BEB.

In a preliminary study on 1 rat, we monitored blinking and spasms of lid closure in a normal rat receiving 7 Hz deep brain stimulation of the subthalamic nucleus 4 hours a day combined with mild dry eye produced by exorbital lacrimal gland removal (36). We tested 3 conditions: 1) 7 Hz subthalamic nucleus deep brain stimulation alone (Fig. 1B, gray bars); 2) 7 Hz subthalamic nucleus deep brain stimulation combined with dry eye (Fig. 1B, black bars); and 3) dry eye alone (Fig. 1B, white bars). In Condition 1, the rat received 5 days of 7 Hz subthalamic nucleus deep brain stimulation alone. In Condition 2, combining the predisposing condition and the environmental trigger, we removed the exorbital gland and the rat received 5 days of 7 Hz subthalamic nucleus deep brain stimulation for 4 hours each day. In Condition 3, we discontinued the 7 Hz subthalamic nucleus deep brain stimulation. For all conditions, we monitored blinking (lid closures <100 milliseconds) and lid spasms (lid closures >100 milliseconds) continuously over a 30-minute period on the last 2 days of each




FIG. 1. An animal model of benign essential blepharospasm using 7 Hz deep brain stimulation (DBS) as the predisposing condition. **A.** A recording of spasms of lid closure and excessive blinking by a rat with dry eye receiving 7 Hz subthalamic nucleus (STN) DBS. **B.** Average number of blinks (# Blinks), blink amplitude (Blink amp), blink duration (Blink Dur), number of spasms (# Spasms), amplitude of spasms (Spasm Amp), and duration of spasms (Spasm Dur) relative to 7 Hz STN DBS alone condition. Spasms were lid closures lasting >100 milliseconds. Error bars are SEM. * $P < 0.05$; *** $P < 0.001$.

condition and normalized all data to the 7 Hz subthalamic nucleus deep brain stimulation alone condition. In the combined 7 Hz subthalamic nucleus deep brain stimulation and dry eye condition, the rat made more blinks than either the 7 Hz subthalamic nucleus deep brain stimulation alone or dry eye alone conditions (Fig. 1B, # Blinks). In the combined 7 Hz subthalamic nucleus deep brain stimulation and dry eye condition, the rat also exhibited more spasms of lid closure than in the other conditions (Fig. 1B, # Spasms). Moreover, the spasm duration was longer in the combined 7 Hz subthalamic nucleus deep brain stimulation and dry eye condition than in the 7 Hz subthalamic nucleus deep brain stimulation alone or dry eye alone condition (Fig. 1A, B, Spasm Dur). Finally, the rat made significantly larger blinks in the combined 7 Hz subthalamic nucleus deep brain stimulation and dry eye condition than in 7 Hz subthalamic nucleus deep brain stimulation alone condition ($P < 0.05$; Fig. 1B, Blink Amp). Although preliminary, these data indicate that the next rodent model of BEB should be developed by combining theta frequency deep brain stimulation of the subthalamic nucleus and dry eye.

Thus far, animal models of BEB have not been tested for the abnormal sensitivity to light associated with BEB (1,24,87). The neural bases of photophobia in patients with BEB are unknown. Physiological and behavioral studies of photophobia implicate changes in blood flow (88), melanopsin ganglion cell inputs to somatosensory thalamic regions (89), intraocular no-

ciceptors (90), and calcitonin gene-related peptide trigeminal sensitization (91,92). Because all of these mechanisms involve elevated trigeminal excitability, we anticipate that rodent models of BEB will also exhibit exaggerated light sensitivity.

The evidence from animal models indicates that spasms of lid closure and trigeminal hyperexcitability of BEB result from exaggerated neuroplasticity, an amplification of the normally adaptive modifications of blinking initiated by eye irritation. The adaptive plasticity initiated by eye irritation seems to involve the cerebellum (29–31), and the exaggeration of plasticity ensues from abnormal basal ganglia modulation of cerebellar activity (82). These results are consistent with the available data pointing to abnormal cerebellar basal ganglia interactions as a major component of dystonia (62,93–96). Although animal models are not identical to human BEB, they are invaluable for identifying the neural mechanisms and circuits causing BEB.

REFERENCES

1. Hallett M, Evinger C, Jankovic J, Stacy M. Update on blepharospasm: report from the BEBRF International Workshop. *Neurology*. 2008;71:1275–1282.
2. Defazio G, Abbruzzese G, Aniello MS, Bloise M, Crisci C, Eleopra R, Fabbrini G, Girlanda P, Liguori R, Macerollo A, Marinelli L, Martino D, Morgante F, Santore L, Tinazzi M, Beradelli A. Environmental risk factors and clinical phenotype in familial and sporadic primary blepharospasm. *Neurology*. 2011;77:631–637.
3. Defazio G, Abbruzzese G, Aniello MS, Di Fede R, Esposito M, Fabbrini G, Girlanda P, Liguori R, Marinelli L, Martino D,

- Morgante F, Santoro L, Tinazzi M, Beradelli A. Eye symptoms in relatives of patients with primary adult-onset dystonia. *Mov Disord.* 2012;27:305–307.
4. **Hallett M.** Blepharospasm: recent advances. *Neurology.* 2002;59:1306–1312.
 5. **Jinnah HA,** Hallett M. In the wink of an eye: nature and nurture in blepharospasm. *Neurology.* 2011;77:616–617.
 6. **Martino D,** Defazio G, Alessio G, Abbruzzese G, Girlanda P, Tinazzi M, Fabbrini G, Marinelli L, Majorana G, Buccafusca M, Vacca L, Livrea P, Beradelli A. Relationship between eye symptoms and blepharospasm: a multicenter case-control study. *Mov Disord.* 2005;20:1564–1570.
 7. **O'Riordan S,** Raymond D, Lynch T, Saunders-Pullman R, Bressman SB, Daly L, Hutchinson M. Age at onset as a factor in determining the phenotype of primary torsion dystonia. *Neurology.* 2004;63:1423–1426.
 8. **Clarimon J,** Brancati F, Peckham E, Valente EM, Dallapiccola B, Abbruzzese G, Girlanda P, Defazio G, Berardelli A, Hallett M, Singleton AB. Assessing the role of DRD5 and DYT1 in two different case-control series with primary blepharospasm. *Mov Disord.* 2007;22:162–166.
 9. **Defazio G,** Brancati F, Valente EM, Caputo V, Pizzuti A, Martino D, Abbruzzese G, Livrea P, Berardelli A, Dallapiccola B. Familial blepharospasm is inherited as an autosomal dominant trait and relates to a novel unassigned gene. *Mov Disord.* 2003;18:207–212.
 10. **Defazio G,** Martino D, Aniello MS, Masi G, Abbruzzese G, Lamberti S, Valente EM, Brancati F, Livrea P, Berardelli A. A family study on primary blepharospasm. *J Neurol Neurosurg Psychiatry.* 2006;77:252–254.
 11. **Misbahuddin A,** Placzek MR, Chaudhuri KR, Wood NW, Bhatia KP, Warner TT. A polymorphism in the dopamine receptor DRD5 is associated with blepharospasm. *Neurology.* 2002;58:124–126.
 12. **Xiromerisiou G,** Dardiotis E, Tsironi EE, Hadjigeorgiou G, Ralli S, Kara E, Petalas A, Tachmitzi S, Hardy J, Houlden H. THAP1 mutations in a Greek primary blepharospasm series. *Parkinsonism Relat Disord.* 2013;19:404–405.
 13. **Defazio G,** Livrea P, Guanti G, Lepore V, Ferrari E. Genetic contribution to idiopathic adult-onset blepharospasm and cranial-cervical dystonia. *Eur Neurol.* 1993;33:345–350.
 14. **Dhaenens CM,** Krystkowiak P, Douay X, Charpentier P, Bele S, Destee A, Sablonniere B. Clinical and genetic evaluation in a French population presenting with primary focal dystonia. *Mov Disord.* 2005;20:822–825.
 15. **Berardelli A,** Rothwell JC, Day BL, Marsden CD. Pathophysiology of blepharospasm and oromandibular dystonia. *Brain.* 1985;108:593–608.
 16. **Jankovic J,** Havins WE, Wilkins RB. Blinking and blepharospasm. Mechanism, diagnosis, and management. *JAMA.* 1982;248:3160–3164.
 17. **Marsden CD.** The focal dystonias. *Clin Neuropharmacol.* 1986;9(suppl 2):S549–S560.
 18. **Patel BC,** Anderson RL. Blepharospasm and related facial movement disorders. *Curr Opin Ophthalmol.* 1995;6:86–899.
 19. **Eekhof JL,** Aramideh M, Speelman JD, Ongerboer de Visser BW. Orbicularis oculi and orbicularis oris reflexes in blepharospasm and torticollis spasmodica during spasm-free intervals. *Eur Neurol.* 2001;46:75–78.
 20. **Evinger C,** Mao JB, Powers AS, Kassem IS, Schicatano EJ, Henriquez VM, Peshori KR. Dry eye, blinking, and blepharospasm. *Mov Disord.* 2002;17(suppl 2):S75–S78.
 21. **Peshori KR,** Schicatano EJ, Gopalaswamy R, Sahay E, Evinger C. Aging of the trigeminal blink system. *Exp Brain Res.* 2001;136:351–363.
 22. **Eekhof JL,** Aramideh M, Bour LJ, Hilgevoord AA, Speelman HD, Ongerboer de Visser BW. Blink reflex recovery curves in blepharospasm, torticollis spasmodica, and hemifacial spasm. *Muscle Nerve.* 1996;19:10–15.
 23. **Bentivoglio AR,** Daniele A, Albanese A, Tonali PA, Fasano A. Analysis of blink rate in patients with blepharospasm. *Mov Disord.* 2006;21:1225–1229.
 24. **Adams WH,** Digre KB, Patel BC, Anderson RL, Warner JE, Katz BJ. The evaluation of light sensitivity in benign essential blepharospasm. *Am J Ophthalmol.* 2006;142:82–87.
 25. **Emoto H,** Suzuki Y, Wakakura M, Horie C, Kiyosawa M, Mochizuki M, Kawasaki K, Oda K, Ishiwata K, Ishii K. Photophobia in essential blepharospasm—a positron emission tomographic study. *Mov Disord.* 2010;25:433–439.
 26. **McCann JD,** Gauthier M, Morschbacher R, Goldberg RA, Anderson RL, Fine PG, Digre KB. A novel mechanism for benign essential blepharospasm. *Ophthal Plast Reconstr Surg.* 1999;15:384–389.
 27. **McCarty CA,** Bansal AK, Livingston PM, Stanislavsky YL, Taylor HR. The epidemiology of dry eye in Melbourne. *Aust Ophthalmol.* 1998;105:1114–1119.
 28. **Tsubota K,** Hata S, Okusawa Y, Egami F, Ohtsuki T, Nakamori K. Quantitative videographic analysis of blinking in normal subjects and patients with dry eye. *Arch Ophthalmol.* 1996;114:715–720.
 29. **Chen FP,** Evinger C. Cerebellar modulation of trigeminal reflex blinks: interpositus neurons. *J Neurosci.* 2006;26:10569–10576.
 30. **Evinger C,** Pellegrini JJ, Manning KA. Adaptive gain modification of the blink reflex. A model system for investigating the physiologic bases of motor learning. *Ann N Y Acad Sci.* 1989;563:87–100.
 31. **Pellegrini JJ,** Evinger C. Role of cerebellum in adaptive modification of reflex blinks. *Learn Mem.* 1997;4:77–87.
 32. **Schicatano EJ,** Mantzouranis J, Peshori KR, Partin J, Evinger C. Lid restraint evokes two types of motor adaptation. *J Neurosci.* 2002;22:569–576.
 33. **Choy EP,** Cho P, Benzie IF, Choy CK. Dry eye and blink rate simulation with a pig eye model. *Optom Vis Sci.* 2008;85:129–134.
 34. **Doane MG.** Interactions of eyelids and tears in corneal wetting and the dynamics of the normal human eyeblink. *Am J Ophthalmol.* 1980;89:507–516.
 35. **Doane MG.** Blinking and the mechanics of the lacrimal drainage system. *Ophthalmology.* 1981;88:844–851.
 36. **Kaminer J,** Powers AS, Horn KG, Hui C, Evinger C. Characterizing the spontaneous blink generator: an animal model. *J Neurosci.* 2011;31:11256–11267.
 37. **Nakamori K,** Odawara M, Nakajima T, Mizutani T, Tsubota K. Blinking is controlled primarily by ocular surface conditions. *Am J Ophthalmol.* 1997;124:24–30.
 38. **Berke A.** Blinking frequency and the thickness of the lipid layer. *Adv Exp Med Biol.* 2002;506:513–516.
 39. **Tsubota K.** Tear dynamics and dry eye. *Prog Retin Eye Res.* 1998;17:565–596.
 40. **Tsubota K,** Nakamori K. Dry eyes and video display terminals. *N Engl J Med.* 1993;328:584.
 41. **Quartarone A,** Rizzo V, Morgante F. Clinical features of dystonia: a pathophysiological revisit. *Curr Opin Neurol.* 2008;21:484–490.
 42. **Peterson DA,** Sejnowski TJ, Poizner H. Convergent evidence for abnormal striatal synaptic plasticity in dystonia. *Neurobiol Dis.* 2010;37:558–573.
 43. **Martella G,** Tassone A, Sciamanna G, Platania P, Cuomo D, Visconti MT, Bonsi P, Cacci E, Biagioli S, Usiello A, Bernardi G, Sharma N, Standert DG, Pisani A. Impairment of bidirectional synaptic plasticity in the striatum of a mouse model of DYT1 dystonia: role of endogenous acetylcholine. *Brain.* 2009;132:2336–2349.
 44. **Quartarone A,** Pisani A. Abnormal plasticity in dystonia: disruption of synaptic homeostasis. *Neurobiol Dis.* 2011;42:162–170.
 45. **Quartarone A,** Bagnato S, Rizzo V, Siebner HR, Dattola V, Scalvetti A, Morgante F, Battaglia F, Romano M, Girlanda P. Abnormal associative plasticity of the human motor cortex in writer's cramp. *Brain.* 2003;126:2586–2596.
 46. **Quartarone A,** Morgante F, Sant'angelo A, Rizzo V, Bagnato S, Terranova C, Siebner HR, Berardelli A, Girlanda P. Abnormal plasticity of sensorimotor circuits extends beyond the affected body part in focal dystonia. *J Neurol Neurosurg Psychiatry.* 2008;79:985–990.

47. **Quartarone A**, Rizzo V, Bagnato S, Morgante F, Sant'Angelo A, Romano M, Crupi D, Girlanda P, Rothwell JC, Siebner HR. Homeostatic-like plasticity of the primary motor hand area is impaired in focal hand dystonia. *Brain*. 2005;128:1943–1950.
48. **Quartarone A**, Siebner HR, Rothwell JC. Task-specific hand dystonia: can too much plasticity be bad for you? *Trends Neurosci*. 2006;29:192–199.
49. **Quartarone A**, Sant'Angelo A, Battaglia F, Bagnato S, Rizzo V, Morgante F, Rothwell JC, Siebner HR, Girlanda P. Enhanced long-term potentiation-like plasticity of the trigeminal blink reflex circuit in blepharospasm. *J Neurosci*. 2006;26:716–721.
50. **Schicatano EJ**, Basso MA, Evinger C. Animal model explains the origins of the cranial dystonia benign essential blepharospasm. *J Neurophysiol*. 1997;77:2842–2846.
51. **Perlmutter JS**, Tempel LW, Black KJ, Parkinson D, Todd RD. MPTP induces dystonia and parkinsonism. Clues to the pathophysiology of dystonia. *Neurology*. 1997;49:1432–1438.
52. **Horie C**, Suzuki Y, Kiyosawa M, Mochizuki M, Wakakura M, Oda K, Ishiwata K, Ishii K. Decreased dopamine D receptor binding in essential blepharospasm. *Acta Neurol Scand*. 2009;119:49–54.
53. **Perlmutter JS**, Stambuk MK, Markham J, Black KJ, McGee-Minnich L, Jankovic J, Moerlein SM. Decreased [¹⁸F]spiperone binding in putamen in idiopathic focal dystonia. *J Neurosci*. 1997;17:843–850.
54. **Campbell DB**, Hess EJ. L-type calcium channels contribute to the tottering mouse dystonic episodes. *Mol Pharmacol*. 1999;55:23–31.
55. **Fan X**, Hughes KE, Jinnah HA, Hess EJ. Selective and sustained AMPA receptor activation in cerebellum induces dystonia in mice. *J Pharmacol Exp Ther*. 2012;340:733–741.
56. **Fan X**, Hughes KE, Jinnah HA, Hess EJ. Selective and sustained alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor activation in cerebellum induces dystonia in mice. *J Pharmacol Exp Ther*. 2012;340:733–741.
57. **Jinnah HA**, Hess EJ. A new twist on the anatomy of dystonia: the basal ganglia and the cerebellum? *Neurology*. 2006;67:1740–1741.
58. **LeDoux MS**. Animal models of dystonia: lessons from a mutant rat. *Neurobiol Dis*. 2011;42:152–161.
59. **LeDoux MS**, Hurst DC, Lorden JF. Single-unit activity of cerebellar nuclear cells in the awake genetically dystonic rat. *Neuroscience*. 1998;86:533–545.
60. **LeDoux MS**, Lorden JF. Abnormal cerebellar output in the genetically dystonic rat. *Adv Neurol*. 1998;78:63–78.
61. **LeDoux MS**, Lorden JF, Ervin JM. Cerebellectomy eliminates the motor syndrome of the genetically dystonic rat. *Exp Neurol*. 1993;120:302–310.
62. **Neychev VK**, Fan X, Mitev VI, Hess EJ, Jinnah HA. The basal ganglia and cerebellum interact in the expression of dystonic movement. *Brain*. 2008;131:2499–2509.
63. **Pizoli CE**, Jinnah HA, Billingsley ML, Hess EJ. Abnormal cerebellar signaling induces dystonia in mice. *J Neurosci*. 2002;22:7825–7833.
64. **Brown P**. Oscillatory nature of human basal ganglia activity: relationship to the pathophysiology of Parkinson's disease. *Mov Disord*. 2003;18:357–363.
65. **de Solages C**, Hill BC, Koop MM, Henderson JM, Bronte-Stewart H. Bilateral symmetry and coherence of subthalamic nuclei beta band activity in Parkinson's disease. *Exp Neurol*. 2010;221:260–266.
66. **Doyle LM**, Kuhn AA, Hariz M, Kupsch A, Schneider GH, Brown P. Levodopa-induced modulation of subthalamic beta oscillations during self-paced movements in patients with Parkinson's disease. *Eur J Neurosci*. 2005;21:1403–1412.
67. **Gatev P**, Darbin O, Wichmann T. Oscillations in the basal ganglia under normal conditions and in movement disorders. *Mov Disord*. 2006;21:1566–1577.
68. **Hammond C**, Bergman H, Brown P. Pathological synchronization in Parkinson's disease: networks, models and treatments. *Trends Neurosci*. 2007;30:357–364.
69. **Jenkinson N**, Brown P. New insights into the relationship between dopamine, beta oscillations and motor function. *Trends Neurosci*. 2011;34:611–618.
70. **Kuhn AA**, Kempf F, Brucke C, Gaynor Doyle L, Martinez-Torres I, Pogosyan A, Trottenberg T, Kupsch A, Schneider GH, Hariz MI, Vandenberghe W, Nuttin B, Brown P. High-frequency stimulation of the subthalamic nucleus suppresses oscillatory beta activity in patients with Parkinson's disease in parallel with improvement in motor performance. *J Neurosci*. 2008;28:6165–6173.
71. **Weinberger M**, Hutchison WD, Alavi M, Hodaie M, Lozano AM, Moro E, Dostrovsky JO. Oscillatory activity in the globus pallidus internus: comparison between Parkinson's disease and dystonia. *Clin Neurophysiol*. 2012;123:358–368.
72. **Chen CC**, Kuhn AA, Trottenberg T, Kupsch A, Schneider GH, Brown P. Neuronal activity in globus pallidus interna can be synchronized to local field potential activity over 3–12 Hz in patients with dystonia. *Exp Neurol*. 2006;202:480–486.
73. **Liu X**, Wang S, Yianni J, Nandi D, Bain PG, Gregory R, Stein JF, Aziz TZ. The sensory and motor representation of synchronized oscillations in the globus pallidus in patients with primary dystonia. *Brain*. 2008;131:1562–1573.
74. **Starr PA**, Rau GM, Davis V, Marks WJ Jr, Ostrem JL, Simmons D, Lindsey N, Turner RS. Spontaneous pallidal neuronal activity in human dystonia: comparison with Parkinson's disease and normal macaque. *J Neurophysiol*. 2005;93:3165–3176.
75. **Brucke C**, Huebl J, Schonecker T, Neumann WJ, Yarrow K, Kupsch A, Blahak C, Lutjens G, Brown P, Krauss JK, Schneider G-H, Kuhn AA. Scaling of movement is related to pallidal gamma oscillations in patients with dystonia. *J Neurosci*. 2012;32:1008–1019.
76. **Kuhn AA**, Doyle L, Pogosyan A, Yarrow K, Kupsch A, Schneider GH, Hariz MI, Trottenberg T, Brown P. Modulation of beta oscillations in the subthalamic area during motor imagery in Parkinson's disease. *Brain*. 2006;129:695–706.
77. **Kuhn AA**, Williams D, Kupsch A, Limousin P, Hariz M, Schneider GH, Yarrow K, Brown P. Event-related beta desynchronization in human subthalamic nucleus correlates with motor performance. *Brain*. 2004;127:735–746.
78. **Leventhal DK**, Gage GJ, Schmidt R, Pettibone JR, Case AC, Berke JD. Basal ganglia beta oscillations accompany cue utilization. *Neuron*. 2012;73:523–536.
79. **Levy R**, Ashby P, Hutchison WD, Lang AE, Lozano AM, Dostrovsky JO. Dependence of subthalamic nucleus oscillations on movement and dopamine in Parkinson's disease. *Brain*. 2002;125:1196–1209.
80. **Priore A**, Foffani G, Pesenti A, Bianchi A, Chiesa V, Baselli G, Caputo E, Tamia F, Rampini P, Egidi M, Locatelli M, Barbieri S, Scarlato G. Movement-related modulation of neural activity in human basal ganglia and its L-DOPA dependency: recordings from deep brain stimulation electrodes in patients with Parkinson's disease. *Neurol Sci*. 2002;23(suppl 2):101–102.
81. **Williams D**, Kuhn A, Kupsch A, Tijssen M, van Bruggen G, Speelman H, Hotton G, Yarrow K, Brown P. Behavioural cues are associated with modulations of synchronous oscillations in the human subthalamic nucleus. *Brain*. 2003;126:1975–1985.
82. **Kaminer J**, Thakur P, Evinger C. Frequency matters: beta-band subthalamic nucleus deep-brain stimulation induces Parkinsonian-like blink abnormalities in normal rats. *Eur J Neurosci*. 2014;40:3237–3742.
83. **Mao JB**, Evinger C. Long-term potentiation of the human blink reflex. *J Neurosci*. 2001;21:RC151.
84. **Ryan M**, Kaminer J, Enmore P, Evinger C. Trigeminal high-frequency stimulation produces short- and long-term modification of reflex blink gain. *J Neurophysiol*. 2014;111:888–895.
85. **Battaglia F**, Ghilardi MF, Quartarone A, Bagnato S, Girlanda P, Hallett M. Impaired long-term potentiation-like plasticity of the trigeminal blink reflex circuit in Parkinson's disease. *Mov Disord*. 2006;21:2230–2233.
86. **Bronstein JM**, Tagliati M, Alterman RL, Lozano AM, Volkmann J, Stefani A, Horak FB, Okun MS, Foote KD, Krack P, Pahwa R, Henderson JE, Hariz MI, Bakay RA, Rezai A, Marks WJ, Moro E, Vitek JL, Weaver FM, Gross RE, DeLong M. Deep brain stimulation for Parkinson disease: an expert consensus and review of key issues. *Arch Neurol*. 2011;68:165.

87. **Evans RW**, Digre KB. Light sensitivity in migraineurs. *Headache*. 2003;43:917–920.
88. **Okamoto K**, Tashiro A, Chang Z, Bereiter DA. Bright light activates a trigeminal nociceptive pathway. *Pain*. 2010;149:235–242.
89. **Noseda R**, Kainz V, Jakubowski M, Gooley JJ, Saper CB, Digre K, Burstein R. A neural mechanism for exacerbation of headache by light. *Nat Neurosci*. 2010;13:239–245.
90. **Dolgonos S**, Ayyala H, Evinger C. Light-induced trigeminal sensitization without central visual pathways: another mechanism for photophobia. *Invest Ophthalmol Vis Sci*. 2011;52:7852–7858.
91. **Kaiser EA**, Kuburas A, Recober A, Russo AF. Modulation of CGRP-induced light aversion in wild-type mice by a 5-HT(1B/D) agonist. *J Neurosci*. 2012;32:15439–15449.
92. **Recober A**, Kuburas A, Zhang Z, Wemmie JA, Anderson MG, Russo AF. Role of calcitonin gene-related peptide in light-aversive behavior: implications for migraine. *J Neurosci*. 2009;29:8798–8804.
93. **Argyelan M**, Carbon M, Niethammer M, Ulug AM, Voss HU, Bressman SB, Dhawan V, Edelberg D. Cerebellothalamic cortical connectivity regulates penetrance in dystonia. *J Neurosci*. 2009;29:9740–9747.
94. **Galardi G**, Perani D, Grassi F, Bressi S, Amadio S, Antoni M, Comi GC, Canal N, Fazio F. Basal ganglia and thalamo-cortical hypermetabolism in patients with spasmodic torticollis. *Acta Neurol Scand*. 1996;94:172–176.
95. **Prudente CN**, Hess EJ, Jinnah HA. Dystonia as a network disorder: what is the role of the cerebellum? *Neuroscience*. 2013;260C:23–35.
96. **Sadnicka A**, Hoffland BS, Bhatia KP, van de Warrenburg BP, Edwards MJ. The cerebellum in dystonia—help or hindrance? *Clin Neurophysiol*. 2012;123:65–70.

Benign Essential Blepharospasm—There Is More to It Than Just Blinking

Kathleen B. Digre, MD

Journal of Neuro-Ophthalmology 2015;35:379–381
doi: 10.1097/WNO.0000000000000316
© 2015 by North American Neuro-Ophthalmology Society

Benign essential blepharospasm (BEB) is recognized today as a primary dystonia causing excessive blinking, squeezing, and involuntary contractions of the orbicularis oculi muscles. This involuntary lid closure leads to functional blindness and decreased quality of life. Besides the blinking and squeezing, patients with BEB are known to have trigeminal hyperexcitability as demonstrated by blink reflex testing and photophobia. Patients with BEB frequently use sensory tricks, like touching the side of the eye, humming, or singing that will temporarily improve the spasms. For decades, this led clinicians to consider blepharospasm to be a nonphysiological disorder. However, many studies in the last 60 years have dispelled that belief.

The condition occurs more frequently in women by a ratio of almost 3 to 1. Most are white. Although the median age is approximately 53 years, blepharospasm occasionally has been reported in children. Many individ-

uals go years before they are appropriately diagnosed. The most valid findings to make the diagnosis are involuntary eyelid narrowing or closure due to spasms of the orbicularis oculi muscle, bilateral spasms that are synchronous and stereotyped, a sensory trick, and inability to suppress the spasms and blink count voluntarily (1). Many individuals report that there is a family history of dystonia or benign tremor or Parkinson disease. Some predisposing factors are believed to be recent stressful events, a history of dry eye or keratitis, and head trauma (2). BEB has profound effects on visual quality of life and overall quality of life, and there is a tendency to more depression (3). For such a disabling condition, we have limited treatment options. There is a real need for greater understanding of this disorder and better treatments to help our patients.

In the accompanying article, Evinger (4) reviews what animal models teach us about this vexing condition. These models provide hope that if we can model a condition in an animal, we are more likely to be able to understand factors that cause it and create more effective treatments for our patients.

Initially, Evinger reminds us that the etiology of BEB may occur due to a predisposition (e.g., genetic) and an environmental trigger—the so called “2 hit” hypothesis. Although there is no known gene for the condition, frequency of a positive family history suggests that there is a genetic component. But there must also be an environmental trigger. Epidemiological data strongly point to the association of dry eyes and blepharitis as potential environmental triggers.

What dry eye and dry eye symptoms do in predisposed individuals is to exaggerate neuroplasticity by increasing blink frequency and amplitude in an attempt to restore tears. Modifying the trigeminal blink reflex becomes

Department of Ophthalmology and Visual Sciences, Moran Eye Center, University of Utah, Salt Lake City, Utah.

Supported by a grant to the Department of Ophthalmology and Visual Sciences from Research to Prevent Blindness, Inc, New York, NY.

K. B. Digre is listed as an inventor on a patent pending for thin-film coatings designed for the treatment of photophobia; she could receive royalties on any commercial sales of these coatings.

Address correspondence to Kathleen B. Digre, MD, Department of Ophthalmology and Visual Sciences, John Moran Eye Center, 65 N Mario Capecchi Drive, Salt Lake City, UT 84132; E-mail: Kathleen.digre@hsc.utah.edu